

U-Pb and Lu-Hf isotopic of zircon from (poly)metamorphic anorthosites and metabasites from the Central Zone of the Limpopo Belt

Zeh, Armin¹ Gerdes, Axel² Klemm, Reiner³

¹Universität Würzburg, Lehrstuhl für Geodynamik and Geomaterialforschung, Am Hubland, D-97074 Würzburg, Germany. ²Universität Frankfurt, Institut für Geowissenschaften, Altenhöfer Allee 1, D-60438 Frankfurt am Main, Germany ³Geoforschungszentrum, Nordbayern, Schlossgarten 5, D-91054 Erlangen, Germany

U-Pb and Lu-Hf isotope – laser ablation ICP(MC)-MS - analyses of zircon grains/domains from two anorthosite samples of the Messina Layered Intrusion (eastern part of the Limpopo Belt) provide evidence for three zircon forming events: at 3.35 Ga, 2.65 Ga and 2.03 Ga. Zircon formation – rather than zircon alteration - during all three events is constrained by increasing $^{176}\text{Hf}/^{177}\text{Hf}$ from ~ 0.2806 (at 3.35 Ga), to ~ 0.2010 (at 2.65 Ga), and ~ 0.2813 (at 2.0 Ga). This increase also indicates that a melt phase was present during zircon growth at all three times (see Gerdes & Zeh, 2008). Thus, the combined U-Pb and Lu-Hf datasets (in combination with CL images) provide evidence that the Messina layered intrusion emplaced at 3.35 Ga, and that their anorthositic differentiation products underwent re-melting during high-grade metamorphic events at 2.65 and 2.03 Ga. These conclusions well conform to those previously obtained from the nearby Sand River Gneiss TTG suite (Gerdes & Zeh, 2008). Epsilon Hf (3.35Ga) between +1.5 and -2.5 of the oldest zircon domains indicates that the Messina layered intrusion was derived from a Paleoproterozoic mantle source, which had a nearly chondritic composition. Notably, chondritic values were also obtained for the second zircon generation formed at 2.65 Ga. These values, however, can be explained by crustal fractionation during the re-melting process.

CL images and U-Pb ages of zircon grains of a metabasic rock from the Venetia area (central part of the Limpopo Belt) also provide evidence for Paleoproterozoic magmatic activities at 3.3 Ga, and for a metamorphic overprint at about 2.0 Ga. Textural relationships, furthermore, hint that at least some of the Paleoproterozoic zircon grains/domains were formed under amphibolite-facies conditions (650°C/6-7 kbar), perhaps as a result of the prograde mineral reaction: Hbl(Zr-rich) + Chl + Grt = Cumingtonit + Pl + Zrc. Presently, Lu-Hf analyses are under progress to test this hypothesis.

References

Gerdes A, Zeh, A (2008) Chemical Geology, doi: 10.1016/j.chemgeo.2008.03.005

Abs. No. **15**
Meeting: **DMG 2008**
submitted by: **Zeh, Armin**
email: **armin.zeh@mail.uni-
wuerzburg.de**
date: **0000-00-00**
Req. presentation: **Vortrag**
Req. session: **S05**