

Barium and Sr isotope compositions in carbonaceous chondrites

Wombacher, Frank¹ Hammerschmidt, Konrad¹ Becker, Harry¹

¹ Institut für Geologische Wissenschaften, Freie Universität Berlin, 12249 Berlin, Germany

To further evaluate the distribution of nucleosynthetic isotope anomalies in bulk solar system samples, precise Ba and Sr isotope compositions were determined by TIMS in chondrites (Murchison CM2, Allende CV3, Kainsaz CO3, Forest City H5, St. Severin LL6) and a HCl leachate from Murchison. Of the bulk samples, only the carbonaceous chondrites display slight enrichments in the r-process dominated nuclides ¹³⁵Ba and ¹³⁷Ba. The HCl leachate from Murchison shows a much larger r-process isotope excess of 249, 136 and 59 ppm for ¹³⁵Ba, ¹³⁷Ba and ¹³⁸Ba, respectively. None of the samples displays resolvable anomalies for the rare p-only nuclides ¹³⁰Ba and ¹³²Ba, outside the external reproducibility of 175 and 104 ppm (2 sd). Stable Sr isotopes in Allende and Murchison (bulk and HCl leachate) display normal compositions within 49 ppm (2 sd; ⁸⁶Sr/⁸⁸Sr rel. to ⁸⁴Sr/⁸⁸Sr). The above Ba and Sr isotope data are in accord with previous data by Hidaka et al. (2003), Carlson et al. (2007) and Andreasen & Sharma (2007).

Discussion and conclusions: Testruns using a container-less laser fusion method (Pack et al. 2007) indicate that the anomalies do not result from incomplete sample digestion. The Murchison HCl leachate shows that the anomalous Ba resides in chemically labile carrier phases and that Sr is either not abundant in these phases or isotopically normal. This is remarkable, since Ba and Sr feature a similar condensation temperature and chemical behaviour and may imply that the production of Sr and Ba r-process isotopes is decoupled. Also, Ba and Sr p-process isotopes are normal and apparently decoupled from Sm isotopes (Andreasen & Sharma 2006). Carbonaceous chondrite leachates display large isotope anomalies in the refractory elements Ba, Mo, Zr, Os that are (nearly) balanced in the bulk samples. This requires that dust in the carbonaceous chondrite source region was never completely vaporized.

References

Andreasen R, Sharma M (2006) Solar nebula heterogeneity in p-process samarium and neodymium isotopes. *Science* 314:806-809.

Andreasen R, Sharma M (2007) Mixing and homogenization in the early solar system: Clues from Sr, Ba, Nd, and Sm isotopes in meteorites. *Astrophys J* 665:874-883.

Carlson RW, Boyet M, Horan M (2007) Chondrite barium, neodymium, and samarium isotopic heterogeneity and early earth differentiation. *Science* 316:1175-1178.

Hidaka H, Ohta Y, Yoneda S (2003) Nucleosynthetic components in the early solar system inferred from Ba isotopic compositions in carbonaceous chondrites. *Earth Planet Sci Lett* 214:455-466.

Pack A, Russell SS, Shelley JMG, van Zuilen M (2007) Geo- and cosmochemistry of the twin elements yttrium and holmium. *Geochim Cosmochim Acta* 71:4592-4608.

Abs. No. **60**
Meeting: **DMG 2008**
submitted by: **Wombacher, Frank**
email: **fwo@zedat.fu-berlin.de**
date: **2008-05-26**
Req. presentation: **Poster**
Req. session: **S02**